

RATIONALLY CONNECTED VARIETIES OVER THE MAXIMALLY UNRAMIFIED EXTENSION OF p -ADIC FIELDS

BRADLEY DUESLER AND AMANDA KNECHT

ABSTRACT. A result of Graber, Harris, and Starr shows that a rationally connected variety defined over the function field of a curve over the complex numbers always has a rational point. Similarly, a separably rationally connected variety over a finite field or the function field of a curve over any algebraically closed field will have a rational point. Here we show that rationally connected varieties over the maximally unramified extension of the p -adics usually, in a precise sense, have rational points. This result is in the spirit of Ax and Kochen's result saying that the p -adics are usually C_2 fields. The method of proof utilizes a construction from mathematical logic called the ultraproduct.

1. INTRODUCTION

Let X be a proper, smooth variety over a field K and \overline{K} an algebraic closure of K . A guiding principle in the study of K -rational points on X is given by Kollár ([Kol96] IV.6.3):

Principle 1. *If $\overline{X} = X \times_{\mathrm{Spec}(K)} \mathrm{Spec}(\overline{K})$ is rationally connected, then X should have lots of K -points, at least if K is nice (e.g. K is a finite field, a function field of a curve, or a sufficiently large number field).*

The term “nice” has since been replaced by many with the term quasi-algebraically closed. A field K is said to be quasi-algebraically closed or C_1 if every homogeneous polynomial over K with degree less than the number of variables has a nontrivial solution in K . Some well known examples are finite fields, function fields in one variable over an algebraically closed field, the field of Laurent series over an algebraically closed field and the maximal unramified extension of p -adic fields, \mathbb{Q}_p^{nr} .

A homogeneous polynomial in n variables over K defines a hypersurface in projective $(n - 1)$ -space \mathbb{P}^{n-1} . The hypersurface associated to a form with degree less than the number of variables, when smooth, is a Fano variety, thus is rationally connected (see [Cam92], [KMM92]). Since smooth rationally connected hypersurfaces defined over quasi-algebraically closed fields always have a K -rational point, it is natural to ask:

Question 2. ([Wit10] 1.11) *Let X be a proper, smooth separably rationally connected variety over a field K where K is a quasi-algebraically closed field. Is $X(K) = \emptyset$?*

Affirmative answers to this question have been given when:

B.D. was supported by NSF grants DMS-0134259 and DMS-0240058 and A.K. was supported by NSF grants DMS-0502170.

The second author was supported in part by NSF Grant DMS-0502170.

- (1) K is the function field of a curve defined over an algebraically closed field of characteristic zero [GHS03].
- (2) K is the function field of a curve defined over an algebraically closed field of positive characteristic [dJS03].
- (3) K is a finite field [Esn03].
- (4) $K = k((t))$ is the field of Laurent series over an algebraically closed field [CT11].
- (5) If X is a smooth, proper, rational *surface* over a quasi-algebraically closed field K , then $X(K) \neq \emptyset$ [Man66], [CT87].

Colliot-Thélène and Madore have shown that there exist fields K of cohomological dimension 1 and del Pezzo surfaces X_d of degrees $d = 2, 3, 4$ such that $X_d(K) = \emptyset$ [CTM04]. In particular, these fields are examples of fields of cohomological dimension 1 which are not C_1 . This seems to rule out a cohomological proof that separably rationally connected varieties over C_1 fields have points.

Let us recall some basic facts about rational connectivity. Suppose that X is a smooth, projective variety defined over an arbitrary field K . We say that X is *separably rationally connected* if there is a variety Y and a morphism $u : Y \times \mathbb{P}^1 \rightarrow X$ such that

$$u^{(2)} : (Y \times \mathbb{P}^1) \times (Y \times \mathbb{P}^1) \rightarrow X \times X$$

is dominant and smooth at the generic point. If the field K is algebraically closed, we can simplify the definition and say that X is *separably rationally connected* if there is a rational curve, called a very free curve, $f : \mathbb{P}^1 \rightarrow X$ such that f^*T_X is ample ([Kol96] IV.3.7). Over an arbitrary field K we say that X is *rationally connected* if there is a family of proper algebraic curves $g : U \rightarrow Y$ whose geometric fibers are irreducible rational curves with cycle morphism $u : U \rightarrow X$ such that $u^{(2)}$ is dominant. If the field K is uncountable and algebraically closed, then we say that X is *rationally connected* if for very general closed points $x_1, x_2 \in X$ there is a morphism $f : \mathbb{P}^1 \rightarrow X$ such that $x_1, x_2 \in f(\mathbb{P}^1)$ ([Kol96] IV.3.6). Over any field of characteristic zero, the notions of rationally connected and separably rationally connected are equivalent ([Kol96] IV.3.3).

The proof that rational connectivity and separable rational connectivity are equivalent over characteristic zero relies in some way on generic smoothness, which fails in characteristic p . In fact, in positive characteristic there are smooth, projective varieties for which one can find a rational curve through any two closed points, but the variety does not contain any very free curves [SK79]. One can think of this as rationally connected varieties containing lots of rigid rational curves while separably rationally connected varieties have rational curves that freely deform. Another example of a rationally connected but not separably rationally connected variety over a field of positive characteristic is give by Kollár ([Kol96] V.5.19).

Because of the subtleties between rationally and separably rationally connected over characteristic p , we had to be careful when stating Question 2. But in this paper we are considering varieties defined over fields of characteristic zero, so the terms are interchangeable. The question we consider in this article is whether or not a smooth, projective, rationally connected variety over the maximal unramified extension of the p -adics, \mathbb{Q}_p^{nr} , has a rational point. Lang's theorem asserts that this is true for Fano hypersurfaces [Lan52]. Here we prove a partial result.

Theorem 3. *Fix a numerical polynomial P . There is a finite set of exceptional primes $e(P)$, depending only on P , so that if X is a smooth, projective, rationally*

connected variety defined over \mathbb{Q}_p^{nr} with Hilbert polynomial P , then $X(\mathbb{Q}_p^{nr}) \neq \emptyset$ as long as $p \notin e(P)$.

A polynomial $P(z) \in \mathbb{Q}(z)$ is called a *numerical polynomial* if $P(n)$ is an integer for all sufficiently large integers n .

This theorem is similar to Ax and Kochen's theorem [AK65] that the p -adic number fields are almost C_2 . Emil Artin conjectured that the p -adic fields \mathbb{Q}_p are C_2 . In general, a C_i field K is one for which any form in $K[x_1, \dots, x_n]_d$ with $n > d^i$ has a nontrivial zero. In [Ter66] Terjanian found a counterexample to Artin's conjecture, see for instance [Ser73]. However, using the methods of mathematical logic, Ax and Kochen were able to show that \mathbb{Q}_p is almost C_2 in the following sense.

Theorem 4. (*Ax and Kochen*) Fix an integer $d > 0$. Then there exists a finite number of primes p_0, \dots, p_m such that for all forms $f \in \mathbb{Q}_p[x_1, \dots, x_n]_d$ with $n > d^2$ and $p \neq p_0, \dots, p_m$, f represents zero over \mathbb{Q}_p .

Their method of proof uses mathematical logic to make precise the analogy that \mathbb{Q}_p is like $\mathbb{F}_p((t))$. Then using the fact that the field $\mathbb{F}_p((t))$ is C_2 [Gre66] is enough for Ax and Kochen to conclude the above theorem. We similarly make an analogy between the asymptotic properties of \mathbb{Q}_p^{nr} when p goes to infinity and the properties of $\mathbb{C}((t))$, the field of Laurent expansions over the complex numbers. Then we use the fact that every rationally connected variety over $\mathbb{C}((t))$ contains a $\mathbb{C}((t))$ -point [CT11].

It should be noted that in a recent paper, Denef proves Theorem 4 using only Algebraic Geometry [Den16].

Acknowledgments: We would like to thank Brendan Hassett for many helpful conversations concerning the topics in this paper. We are also grateful to Jean-Louis Colliot-Thélène, Keith Conrad, Olivier Wittenberg, and the referee for their insightful comments.

2. MODEL THEORY AND ALGEBRAIC GEOMETRY

The main tool from Model Theory we will use is the ultraproduct. A more thorough introduction to ultrafilters and ultraproducts is given in [Koc75].

Definition 5. Let S be a set and let Σ be a collection of non-empty subsets of S . Then Σ is called a *non-principal filter* if the following hold:

- (1) $S_1, S_2 \in \Sigma$ implies $S_1 \cap S_2 \in \Sigma$
- (2) $S_1 \in \Sigma$ and $S_2 \supset S_1$ implies $S_2 \in \Sigma$
- (3) For each $s \in S$ there is a set $S_1 \in \Sigma$ such that $s \notin S_1$.

Σ is called a *non-principal ultrafilter* if it is maximal among the class of all non-principal filters on S , or equivalently:

- (4) $S_1 \notin \Sigma$ implies $S - S_1 \in \Sigma$.

Conditions 1, 2, and 4 define an *ultrafilter*.

A simple, but important property of ultrafilters to keep in mind is that if S is the disjoint union of subsets S_1, \dots, S_n , then precisely one of these subsets is in Σ . This observation follows from properties 1 and 4. Namely, at least one of the S_i is in Σ by property 4. Moreover, two disjoint subsets cannot both be in Σ since then so would their intersection, but Σ consists only of nonempty subsets of S .

Given any subset $S_0 \subset S$ it will be useful to know if we can find a non-principal ultrafilter on S containing S_0 . Certainly, if S_0 is a finite set, then properties 3 and 4 of the definition above will prevent us from finding a non-principal ultrafilter containing S_0 . However, this is the only obstruction as the lemma below asserts.

Lemma 6. *Given any infinite subset $S_0 \subset S$, there exists a non-principal ultrafilter containing S_0 .*

Proof. Let Σ consist of all the subsets of S that contain all but a finite number of points in S_0 . It is easy to check that Σ is a non-principal filter on S containing S_0 . The non-principal ultrafilter desired is any maximal filter containing Σ . \square

Our usage of ultrafilters will be for an auxiliary construction called the ultraproduct. In particular, given a collection of fields indexed by a set S , and an ultrafilter Σ on S , we will construct a new field via the ultraproduct. We will use a similar construction for modules.

Definition 7. Given an ultrafilter Σ on S and a collection of rings $\{R_i\}_{i \in S}$ we can form a new ring denoted

$$\prod_{i \in S} R_i / \Sigma$$

defined by componentwise addition and multiplication under the equivalence condition that $a, b \in \prod_{i \in S} R_i$ are equivalent if they agree on a set of indices in Σ . This new ring is called the *ultraproduct* of the R_i 's with respect to Σ . The same definition can be made for groups, modules, etc.

One of the nice aspects of ultraproducts is that the ultraproduct of fields is also a field. Moreover, statements in the language of fields can be transferred between the ultraproduct and its components which leads to the fundamental property of ultraproducts. Let $\{k_p\}_{p \in S}$ be a collection of fields. Then Loš's Theorem ([FJ08] 7.7.1) applied to the particular cases of an ultraproduct of fields can be stated as:

Theorem 8. (Loš) *A first-order formula in the language of rings is true in the ultraproduct of fields $\prod_{p \in S} k_p / \Sigma$ if and only if the set of indices p such that the formula is true in the field k_p is a member of Σ .*

Intuitively, a first-order formula is a formula that only quantifies over elements of the field, not over subsets, sets of subsets, etc. To get a feeling for why Loš's theorem is true, even for structures more general than fields, consider the following lemma.

Lemma 9. *Let N be a positive integer. For each $i \in S$, let M_i be a free module of rank less than N over a ring R_i . Then an ultraproduct of the M_i 's is a free module of rank less than N over the corresponding ultraproduct of the R_i 's.*

Proof. First, assume that the rank of the M_i 's are all $m > 0$. Then for any ultrafilter Σ on S

$$M := \prod_{i \in S} M_i / \Sigma$$

will be a free module of rank m over

$$R := \prod_{i \in S} R_i / \Sigma.$$

To see this, let e_{i1}, \dots, e_{im} be an R_i basis for M_i . Then note that M has basis $(e_{i1})_{i \in S}, \dots, (e_{im})_{i \in S}$ over R .

Now generally, consider the subsets $S_k \in S$ consisting of those $i \in S$ such that the rank of M_i is k . Then S is the disjoint union of S_1, \dots, S_N . By the remarks on the definition of ultrafilter, there is only one such subset contained in Σ , say $S_m \in \Sigma$. It follows that M has rank m over R . \square

What makes this lemma work is the boundedness of the statement (that the rank is less than N). The similar statement that the ultraproduct of finite rank free modules is finite rank is actually false (say if the rank of the free modules keeps increasing). Loš's theorem does not apply to such a statement because it is not a first-order statement.

Next, we develop some basic algebraic geometry over a general ultraproduct of fields $F = \prod_{i \in S} F_i / \Sigma$ of characteristic zero. Suppose we are given a scheme X of finite type over F . There is a natural process to obtain schemes X_i of finite type over F_i , and for almost every $i \in S$, X_i is nicely related to X . However, the X_i are not unique.

Let's assume first that X is an affine scheme corresponding to the F -algebra

$$F[x_0, \dots, x_n] / I(X).$$

Suppose that f_1, \dots, f_k are generators for $I(X)$. We may write each generator as

$$f_j = \sum a_{j,I} x^I, \quad I \in \mathbb{N}^{n+1}.$$

Let $(a_{j,I}^i)_{i \in S} \in \prod_{i \in S} F_i$ be a representative for $a_{j,I}$. Setting

$$f_j^i = \sum a_{j,I}^i x^I,$$

we define X_i as the affine scheme associated to the ideal generated by the f_j^i . These schemes are not unique as they depend on the choice of representatives for the $a_{j,I}$.

We can perform a similar construction in reverse. Namely given schemes X_i defined over F_i for each $i \in S$, we can define their ultraproduct $X = \prod_{i \in S} X_i / \Sigma$ by taking the f_j^i and lifting them to $f \in F[x_0, \dots, x_n]$. This new object is not necessarily pretty, for example if the degrees of the f_j^i are not bounded. However, we will see that under certain circumstances the ultraproduct is a scheme of finite type and other nice properties of the X_i 's will be inherited by X .

Let $X \subseteq \mathbb{P}^n$ be a projective variety defined over a field F with homogeneous ideal $J(X)$, and let $S(X) = F[x_0, \dots, x_n] / J(X)$ denote its homogeneous coordinate ring. For each integer ℓ , we define the Hilbert function φ_X of X by,

$$\varphi_X(\ell) = \dim_F S(X)_\ell.$$

It is a theorem of Hilbert and Serre ([Har77] I.7.5) that there exists a unique numerical polynomial $P(z) \in \mathbb{Q}[z]$ such that $\varphi_X(\ell) = P(\ell)$ for all $\ell \gg 0$. By definition, the degree of the Hilbert polynomial is the dimension of the variety X . Chardin and Moreno-Socías characterize, in terms of their coefficients, which numerical polynomials are Hilbert polynomials of some projective scheme [CMS03].

Lemma 10. *Given a collection of projective varieties $X_i \subset \mathbb{P}_{F_i}^n$ all with Hilbert polynomial P , the ultraproduct X is a projective variety in \mathbb{P}_F^n with Hilbert polynomial P .*

Proof. Let $J_i \subset F_i[x_0, \dots, x_n]$ be the homogeneous ideal of $X_i \subset \mathbb{P}_{F_i}^n$. For each degree $d > 0$ consider the F_i -vector space $J_{i,d}$ of the homogeneous polynomials of degree d in J_i .

Now define

$$J_d := \prod_{i \in S} J_{i,d}/\Sigma.$$

It is a property of the Hilbert polynomial that for sufficiently large d the rank of $J_{i,d}$ is the same for each $i \in S$, i.e. the Hilbert functions of the X_i are equal for sufficiently large d ([Kol96] I.1.5). Then, the proof of Lemma 9 shows that for $d \gg 0$ the rank of J_d equals the rank of $J_{i,d}$. This yields a homogeneous ideal

$$J := \bigoplus_{d>0} J_d \subset F[x_0, \dots, x_n].$$

The corresponding projective variety X denoted by

$$X := \prod_{i \in S} X_i/\Sigma$$

has Hilbert Polynomial P . □

Other nice results on properties of varieties preserved under the ultraproduct can be found in papers of Arapura [Ara11] and Schoutens [Sch05].

3. PROOF OF THE MAIN THEOREM

Now that we have established some basic knowledge of ultraproducts we can prove the main theorem of this paper, Theorem 3, in a way similar to Ax and Kochen's proof of Theorem 4. First we need to recall a theorem of Ax-Kochen and Eršov [AK65, Erš65].

Theorem 11. (*Ax-Kochen and Eršov*) *Let K and K' be two Henselian valued fields of residual characteristic zero. Assume their residue fields k and k' and their value groups Γ and Γ' are elementary equivalent, that is, they have the same set of true sentences in the language of rings, respectively ordered abelian groups. Then K and K' are elementary equivalent, that is, they satisfy the same set of formulas in the language of valued fields.*

Proof. Proof of Theorem 3.

Fix a numerical polynomial P . Let \mathbb{Q}_p^{nr} denote the maximally unramified extension of the p -adics and let S be the set of all primes. Suppose by way of contradiction that there is an infinite subset of primes $e(P) \subset S$ such that for each $p \in e(P)$ there is a smooth, projective, rationally connected variety $X_p \in \mathbb{P}^n$ defined over \mathbb{Q}_p^{nr} having Hilbert polynomial P and $X(\mathbb{Q}_p^{nr}) = \emptyset$. Now by Lemma 6 there is a non-principal ultrafilter Σ containing $e(P)$, and we can define the non-principal ultraproduct

$$K = \prod_{p \in S} \mathbb{Q}_p^{nr}/\Sigma.$$

Both K and $\mathbb{C}((t))$ are Henselian valued fields of characteristic zero. The residue field of $\mathbb{C}((t))$ is simply \mathbb{C} , so algebraically closed of characteristic zero. The residue field of K is a non-principal ultraproduct of the algebraic closures of the finite fields \mathbb{F}_p and is known to be algebraically closed of characteristic zero ([AK65], Lemma 4). Thus, the residue fields of K and $\mathbb{C}((t))$ are elementary equivalent. Note that

the lemma of Ax and Kochen requires the ultrafilter to be non-principal which is why we need $e(P)$ to be infinite. K has value group a non-principal ultraproduct of the integers, otherwise known as an ultrapower of \mathbb{Z} , hence is elementary equivalent to \mathbb{Z} , the value group of $\mathbb{C}((t))$. Thus by Theorem 11, K and $\mathbb{C}((t))$ are elementary equivalent.

For the numerical polynomial P fixed above, consider all varieties $X \subset \mathbb{P}_k^n$ with Hilbert polynomial P . It is possible to choose a uniform $M \gg 0$ such that the Hilbert function $\varphi_X(M)$ is equal to $P(M)$ for all varieties with Hilbert polynomial P ([Has07] 12.47). Let Gr denote the Grassmannian of codimension- $P(M)$ subspaces of the space of polynomials of degree M in $n + 1$ variables:

$$Gr := Grass \left(\binom{n+M}{M} - P(M), k[x_0, \dots, x_n]_M \right).$$

Let $J(X)_M \subset k[x_0, \dots, x_n]_M$ denote the degree M polynomials vanishing on X . Then, $J(X)_M$ defines a point in the Grassmannian Gr and the set of all projective varieties with Hilbert polynomial P is parameterized by a projective variety known as the Hilbert scheme, $Hilb_P \subset Gr$. Let g denote the smooth morphism that injects $Hilb_P$ into Gr , $g : Hilb_P \hookrightarrow Gr$. Notice that since P is a numerical polynomial, everything here is defined over \mathbb{Q} . Over any field L of characteristic 0, every projective L -variety X with Hilbert polynomial P is realized as a fiber g_b of g above a point $b = J(X)_M \in Gr$. Thus, X has an L -point if and only if g_b has an L -point.

If the fiber g_b is rationally connected for some $b \in Gr$, then there is an open neighborhood $b \in U \subset Gr$ such that the fiber g_u is rationally connected if $u \in U$ ([Kol96] IV.3.11). Thus, over \mathbb{Q} , the set Z of points $b \in Gr$ such that the fiber g_b is rationally connected is an open subset of Gr . Since Gr is a variety and Z is an open subset of Gr , over any field L of characteristic zero $Z(L)$ is definable in the ring language. Furthermore, that definition is actually the same one given over \mathbb{Q} . Also, for any field L of characteristic zero and any rationally connected variety X over L with Hilbert polynomial P , there is an L -point $z \in Z(L)$ such that $X = g_z$ and $X(L) \neq \emptyset \iff g_z(L) \neq \emptyset$.

Now let H_L be the set of points $b \in Gr(L)$ such that the fiber over b is rationally connected and contains an L -point:

$$H_L = \{b \in Gr(L) : b \in Z(L), g_b(L) \neq \emptyset\}.$$

Then $H_L \subset Z(L)$ is a definable subset of $Gr(L)$ defined by the following formula over \mathbb{Q} with an existential quantifier: $b \in H_L$ if and only if the formula “ $b \in Z$ and $\exists x \in Hilb_P$ such that $g_P(x) = b$ ” is true in the field L . This formula does not depend on the field L , but only on the polynomial P .

The result of Colliot-Thélène [CT11] $\mathbb{C}((t))$ tells us that the formula “ $b \in Z$ and $\exists x \in A_P$ such that $g_P(x) = b$ ” is true over $\mathbb{C}((t))$. Then by elementary equivalence it is true over the ultraproduct K . But Loš’s Theorem tells us that this statement is false over K by the way we constructed our non-principal ultrafilter Σ . Thus we arrive at our contradiction, and we have shown that the set of primes $e(P) \subset S$ such that for each $p \in e(P)$ there is a smooth, projective, rationally connected variety $X_p \in \mathbb{P}^n$ defined over \mathbb{Q}_p^{nr} having Hilbert polynomial P and $X(\mathbb{Q}_p^{nr}) = \emptyset$ is finite. \square

REFERENCES

- [AK65] James Ax and Simon Kochen. Diophantine problems over local fields. I. *Amer. J. Math.*, 87:605–630, 1965.
- [Ara11] Donu Arapura. Frobenius amplitude, ultraproducts, and vanishing on singular spaces. *Illinois J. Math.*, 55(4):1367–1384 (2013), 2011.
- [Cam92] F. Campana. Connexité rationnelle des variétés de Fano. *Ann. Sci. École Norm. Sup. (4)*, 25(5):539–545, 1992.
- [CMS03] Marc Chardin and Guillermo Moreno-Socías. Regularity of lex-segment ideals: some closed formulas and applications. *Proc. Amer. Math. Soc.*, 131(4):1093–1102 (electronic), 2003.
- [CT87] Jean-Louis Colliot-Thélène. Arithmétique des variétés rationnelles et problèmes birationnels. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, pages 641–653, Providence, RI, 1987. Amer. Math. Soc.
- [CT11] Jean-Louis Colliot-Thélène. Variétés presque rationnelles, leurs points rationnels et leurs dégénérescences. In *Arithmetic geometry*, volume 2009 of *Lecture Notes in Math.*, pages 1–44. Springer, Berlin, 2011.
- [CTM04] Jean-Louis Colliot-Thélène and David A. Madore. Surfaces de del Pezzo sans point rationnel sur un corps de dimension cohomologique un. *J. Inst. Math. Jussieu*, 3(1):1–16, 2004.
- [Den16] Jan Denef. Geometric proofs of theorems of Ax-Kochen and Eršov. *Amer. J. Math.*, 138(1):181–199, 2016.
- [dJS03] A. J. de Jong and J. Starr. Every rationally connected variety over the function field of a curve has a rational point. *Amer. J. Math.*, 125(3):567–580, 2003.
- [Erš65] Ju. L. Eršov. On elementary theory of maximal normalized fields. *Algebra i Logika Sem.*, 4(3):31–70, 1965.
- [Esn03] Hélène Esnault. Varieties over a finite field with trivial Chow group of 0-cycles have a rational point. *Invent. Math.*, 151(1):187–191, 2003.
- [FJ08] Michael D. Fried and Moshe Jarden. *Field arithmetic*, volume 11 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, third edition, 2008. Revised by Jarden.
- [GHS03] Tom Graber, Joe Harris, and Jason Starr. Families of rationally connected varieties. *J. Amer. Math. Soc.*, 16(1):57–67 (electronic), 2003.
- [Gre66] Marvin J. Greenberg. Rational points in Henselian discrete valuation rings. *Bull. Amer. Math. Soc.*, 72:713–714, 1966.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Has07] Brendan Hassett. *Introduction to algebraic geometry*. Cambridge University Press, Cambridge, 2007.
- [KMM92] János Kollár, Yoichi Miyaoka, and Shigefumi Mori. Rational connectedness and boundedness of Fano manifolds. *J. Differential Geom.*, 36(3):765–779, 1992.
- [Koc75] Simon Kochen. The model theory of local fields. In *1xsy) ISILC Logic Conference (Proc. Internat. Summer Inst. and Logic Colloq., Kiel, 1974)*, pages 384–425. Lecture Notes in Math., Vol. 499. Springer, Berlin, 1975.
- [Kol96] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996.
- [Lan52] Serge Lang. On quasi algebraic closure. *Ann. of Math. (2)*, 55:373–390, 1952.
- [Man66] Ju. I. Manin. Rational surfaces over perfect fields. *Inst. Hautes Études Sci. Publ. Math.*, (30):55–113, 1966.
- [Sch05] Hans Schoutens. Log-terminal singularities and vanishing theorems via non-standard tight closure. *J. Algebraic Geom.*, 14(2):357–390, 2005.
- [Ser73] J.-P. Serre. *A course in arithmetic*. Springer-Verlag, New York, 1973. Translated from the French, Graduate Texts in Mathematics, No. 7.

- [SK79] Tetsuji Shioda and Toshiyuki Katsura. On Fermat varieties. *Tôhoku Math. J. (2)*, 31(1):97–115, 1979.
- [Ter66] Guy Terjanian. Un contre-exemple à une conjecture d'Artin. *C. R. Acad. Sci. Paris Sér. A-B*, 262:A612, 1966.
- [Wit10] Olivier Wittenberg. La connexité rationnelle en arithmétique. In *Variétés rationnellement connexes: aspects géométriques et arithmétiques*, volume 31 of *Panor. Synthèses*, pages 61–114. Soc. Math. France, Paris, 2010.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TX 77005

Current address: Voya Financial, Malvern, PA 19355

E-mail address: bradley.duesler@voya.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, VILLANOVA UNIVERSITY, VILLANOVA, PA 19085

E-mail address: Amanda.Knecht@villanova.edu